

EHRHART POLYNOMIALS OF 3-DIMENSIONAL SIMPLE INTEGRAL CONVEX POLYTOPES

YUSUKE SUYAMA

ABSTRACT. We give an explicit formula on the Ehrhart polynomial of a 3-dimensional simple integral convex polytope by using toric geometry.

1. INTRODUCTION

Let $P \subset \mathbb{R}^d$ be an integral convex polytope of dimension d , that is, a convex polytope whose vertices have integer coordinates. For a non-negative integer l , we write $lP = \{lx \mid x \in P\}$. Ehrhart [2] proved that the number of lattice points in lP can be expressed by a polynomial in l of degree d :

$$|(lP) \cap \mathbb{Z}^d| = c_d l^d + c_{d-1} l^{d-1} + \cdots + c_0.$$

This polynomial is called the *Ehrhart polynomial* of P . It is known that:

- (1) $c_0 = 1$.
- (2) c_{d-1} is half of the sum of relative volumes of facets of P ([1, Theorem 5.6]).
- (3) c_d is the volume of P ([1, Corollary 3.20]).

However, we have no formula on other coefficients of Ehrhart polynomials. In particular, we do not know a formula on c_1 for a general 3-dimensional integral convex polytope. In this paper, we find an explicit formula on c_1 of the Ehrhart polynomial of a 3-dimensional *simple* integral convex polytope, see Theorem 5.

Pommersheim [4] gave a method for computing the $(d-2)$ -nd coefficient of the Ehrhart polynomial of a d -dimensional simple integral convex polytope P by using toric geometry. He obtained an explicit description of the Ehrhart polynomial of a tetrahedron by using this method. Our formula is obtained by using this method for a general 3-dimensional simple integral convex polytope.

The structure of the paper is as follows. In Section 2, we state the main theorem and give a few examples. In Section 3, we give a proof of the main theorem.

ACKNOWLEDGMENT. This work was supported by Grant-in-Aid for JSPS Fellows 15J01000. The author wishes to thank his supervisor, Professor Mikiya Masuda, for his continuing support.

2. THE MAIN THEOREM

Let $P \subset \mathbb{R}^3$ be a 3-dimensional simple integral convex polytope, and let $F_1, \dots, F_n$ be the facets of P . For $k = 1, \dots, n$, we denote by $v_k \in \mathbb{Z}^3$ the inward-pointing primitive normal vector of F_k . For an edge E of P , we denote by $\text{Vol}(E)$ the relative

Date: May 17, 2016.

2010 Mathematics Subject Classification. Primary 52B20; Secondary 52B10, 14M25.

Key words and phrases. integral convex polytopes, Ehrhart polynomials, toric geometry.

volume of E , that is, the length of E measured with respect to the lattice of rank one in the line containing E .

DEFINITION 1. For each edge $E = F_{k_1} \cap F_{k_2}$ of P , we define an integer $m(E)$ and a rational number $s(E)$ as follows:

- (1) We define $m(E) = |((\mathbb{R}v_{k_1} + \mathbb{R}v_{k_2}) \cap \mathbb{Z}^3) / (\mathbb{Z}v_{k_1} + \mathbb{Z}v_{k_2})|$.
- (2) There exists a basis e_1, e_2 for $(\mathbb{R}v_{k_1} + \mathbb{R}v_{k_2}) \cap \mathbb{Z}^3$ such that $v_{k_1} = e_1$ and $v_{k_2} = pe_1 + qe_2$ for some $q > p \geq 0$. Then we define $s(E) = s(p, q)$, where $s(p, q)$ is the Dedekind sum, which is defined by

$$s(p, q) = \sum_{i=1}^q \left(\left(\frac{i}{q} \right) \right) \left(\left(\frac{pi}{q} \right) \right), \quad ((x)) = \begin{cases} x - [x] - \frac{1}{2} & (x \notin \mathbb{Z}), \\ 0 & (x \in \mathbb{Z}). \end{cases}$$

REMARK 2. We have $q = m(E)$. Although p is not uniquely determined, $s(p, q)$ does not depend on the choice of e_1, e_2 . Thus $s(E)$ is well-defined.

DEFINITION 3. For each facet F of P , we define a rational number $C(F)$ as follows. We name vertices and facets around F as in Figure 1. We denote by $v \in \mathbb{Z}^3$ the inward-pointing primitive normal vector of F .




FIGURE 1. vertices and facets around F .

For $i = 1, \dots, r$, we define

$$\varepsilon_i = \det(v, v_{k_{i+1}}, v_{k_i}) > 0, \quad a_i = \frac{\langle \overrightarrow{P_{i-1}Q_{i-1}}, v_{k_{i+1}} \rangle}{\varepsilon_i \langle \overrightarrow{P_{i-1}Q_{i-1}}, v \rangle}, \quad b_i = \frac{\langle \overrightarrow{P_iP_{i+1}}, v_{k_{i-1}} \rangle}{\varepsilon_{i-1} \langle \overrightarrow{P_iP_{i+1}}, v \rangle},$$

where $v_{k_0} = v_{k_r}, v_{k_{r+1}} = v_{k_1}, \varepsilon_0 = \varepsilon_r, P_0 = P_r, P_{r+1} = P_1, Q_0 = Q_r$ and $\langle \cdot, \cdot \rangle$ is the standard inner product on $\mathbb{R}^3$. Then we define

$$C(F) = - \sum_{2 \leq i < j \leq r} a_i \begin{vmatrix} b_{i+1} & \varepsilon_{i+1}^{-1} & 0 & \cdots & 0 \\ \varepsilon_{i+1}^{-1} & b_{i+2} & \varepsilon_{i+2}^{-1} & \ddots & \vdots \\ 0 & \varepsilon_{i+2}^{-1} & \ddots & \ddots & 0 \\ \vdots & \ddots & \ddots & b_{j-2} & \varepsilon_{j-2}^{-1} \\ 0 & \cdots & 0 & \varepsilon_{j-2}^{-1} & b_{j-1} \end{vmatrix} \varepsilon_i \varepsilon_{i+1} \cdots \varepsilon_{j-1} \frac{\text{Vol}(P_{j-1}P_j)}{m(P_{j-1}P_j)},$$

where $P_{j-1}P_j$ is the edge whose endpoints are P_{j-1} and P_j , and the determinants above are understood to be one when $j = i + 1$.

REMARK 4. The proof of Theorem 5 below shows that $C(F)$ does not depend on the choice of F_{k_1} .

The following is our main theorem:

Theorem 5. *Let $P \subset \mathbb{R}^3$ be a 3-dimensional simple integral convex polytope, and let $E_1, \dots, E_m$ and $F_1, \dots, F_n$ be the edges and the facets of P , respectively. Then the coefficient c_1 of the Ehrhart polynomial $|(lP) \cap \mathbb{Z}^3| = c_3 l^3 + c_2 l^2 + c_1 l + c_0$ is given by*

$$\sum_{j=1}^m \left(s(E_j) + \frac{1}{4} \right) \text{Vol}(E_j) + \frac{1}{12} \sum_{k=1}^n C(F_k).$$

EXAMPLE 6. Let a, b, c be positive integers with $\gcd(a, b, c) = 1$ and let $P \subset \mathbb{R}^3$ be the tetrahedron with vertices

$$O = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}, \quad P_1 = \begin{pmatrix} a \\ 0 \\ 0 \end{pmatrix}, \quad P_2 = \begin{pmatrix} 0 \\ b \\ 0 \end{pmatrix}, \quad P_3 = \begin{pmatrix} 0 \\ 0 \\ c \end{pmatrix}.$$

We put $A = \gcd(b, c), B = \gcd(a, c), C = \gcd(a, b)$ and $d = ABC$. Then we have the following table:

edge E	OP_1	OP_2	OP_3	P_1P_2	P_1P_3	P_2P_3
$\text{Vol}(E)$	a	b	c	C	B	A
$m(E)$	1	1	1	cC/d	bB/d	aA/d
$s(E)$	0	0	0	$-s\left(\frac{ab}{d}, \frac{cC}{d}\right)$	$-s\left(\frac{ac}{d}, \frac{bB}{d}\right)$	$-s\left(\frac{bc}{d}, \frac{aA}{d}\right)$

facet F	OP_1P_2	OP_1P_3	OP_2P_3	$P_1P_2P_3$
inward-pointing primitive normal vector of F	$\begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$	$\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$	$\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$	$\begin{pmatrix} -bc/d \\ -ac/d \\ -ab/d \end{pmatrix}$
$C(F)$	ab/c	ac/b	bc/a	$d^2/(abc)$

TABLE 1. the values of $\text{Vol}(E), s(E)$ and $C(F)$.

Thus we have

$$\begin{aligned} & \sum_{E:\text{edge}} \left(s(E) + \frac{1}{4} \right) \text{Vol}(E) + \frac{1}{12} \sum_{F:\text{facet}} C(F) \\ &= \frac{a}{4} + \frac{b}{4} + \frac{c}{4} + \left(-s \left(\frac{ab}{d}, \frac{cC}{d} \right) + \frac{1}{4} \right) C + \left(-s \left(\frac{ac}{d}, \frac{bB}{d} \right) + \frac{1}{4} \right) B \\ &+ \left(-s \left(\frac{bc}{d}, \frac{aA}{d} \right) + \frac{1}{4} \right) A + \frac{1}{12} \left(\frac{ab}{c} + \frac{ac}{b} + \frac{bc}{a} + \frac{d^2}{abc} \right), \end{aligned}$$

which coincides with the formula in [4, Theorem 5].

EXAMPLE 7. Let a and c be positive integers and b be a non-negative integer. Consider the convex hull $P \subset \mathbb{R}^3$ of the six points

$$\begin{aligned} O &= \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}, \quad A = \begin{pmatrix} a \\ 0 \\ 0 \end{pmatrix}, \quad B = \begin{pmatrix} 0 \\ a \\ 0 \end{pmatrix}, \\ O' &= \begin{pmatrix} b \\ 0 \\ c \end{pmatrix}, \quad A' = \begin{pmatrix} a+b \\ 0 \\ c \end{pmatrix}, \quad B' = \begin{pmatrix} b \\ a \\ c \end{pmatrix}. \end{aligned}$$

P is a 3-dimensional simple polytope. We put $g = \gcd(b, c)$. Then we have the following table:

edge E	OA	OB	AB	OO'	AA'	BB'	$O'A'$	$O'B'$	$A'B'$
$\text{Vol}(E)$	a	a	a	g	g	g	a	a	a
$m(E)$	1	c/g	c/g	1	1	c/g	1	c/g	c/g
$s(E)$	0	$-s\left(\frac{b}{g}, \frac{c}{g}\right)$	$s\left(\frac{b}{g}, \frac{c}{g}\right)$	0	0	$-s\left(1, \frac{c}{g}\right)$	0	$s\left(\frac{b}{g}, \frac{c}{g}\right)$	$-s\left(\frac{b}{g}, \frac{c}{g}\right)$

facet F	OAB	$OAA'O'$	$OBB'O'$	$ABB'A'$	$O'A'B'$
inward-pointing primitive normal vector of F	$\begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$	$\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$	$\begin{pmatrix} c/g \\ 0 \\ -b/g \end{pmatrix}$	$\begin{pmatrix} -c/g \\ -c/g \\ b/g \end{pmatrix}$	$\begin{pmatrix} 0 \\ 0 \\ -1 \end{pmatrix}$
$C(F)$	0	c	g^2/c	g^2/c	0

TABLE 2. the values of $\text{Vol}(E)$, $s(E)$ and $C(F)$.




FIGURE 2. the simple polytope P .

Thus we have

$$\begin{aligned}
& \sum_{E:\text{edge}} \left(s(E) + \frac{1}{4} \right) \text{Vol}(E) + \frac{1}{12} \sum_{F:\text{facet}} C(F) \\
&= -s \left(1, \frac{c}{g} \right) g + \frac{3a}{2} + \frac{3g}{4} + \frac{1}{12} \left(c + \frac{2g^2}{c} \right) \\
&= -g \sum_{i=1}^{c/g-1} \left(\frac{i}{g} - \frac{1}{2} \right)^2 + \frac{3a}{2} + \frac{3g}{4} + \frac{c}{12} + \frac{g^2}{6c} \\
&= -g \sum_{i=1}^{c/g-1} \left(\frac{g^2}{c^2} i^2 - \frac{g}{c} i + \frac{1}{4} \right) + \frac{3a}{2} + \frac{3g}{4} + \frac{c}{12} + \frac{g^2}{6c} \\
&= -\frac{g^3}{c^2} \frac{\left(\frac{c}{g} - 1 \right) \frac{c}{g} \left(\frac{2c}{g} - 1 \right)}{6} + \frac{g^2}{c} \frac{\left(\frac{c}{g} - 1 \right) \frac{c}{g}}{2} - g \frac{\frac{c}{g} - 1}{4} + \frac{3a}{2} + \frac{3g}{4} + \frac{c}{12} + \frac{g^2}{6c} \\
&= \frac{3a}{2} + g.
\end{aligned}$$

On the other hand, since

$$\#\{(x, y) \in \mathbb{Z}^2 \mid (x, y, z) \in lP\} = \begin{cases} \frac{(al+1)(al+2)}{2} & ((c/g) \mid z), \\ \frac{al(al+1)}{2} & ((c/g) \nmid z) \end{cases}$$

for $z = 0, 1, \dots, cl$, we have

$$\begin{aligned}
|(lP) \cap \mathbb{Z}^3| &= \frac{(al+1)(al+2)}{2} (gl+1) + \frac{al(al+1)}{2} ((cl+1) - (gl+1)) \\
&= \frac{a^2c}{2} l^3 + \frac{1}{2} (a^2 + ac + 2ag) l^2 + \left(\frac{3a}{2} + g \right) l + 1.
\end{aligned}$$

The coefficient of l is also $3a/2 + g$.

3. PROOF OF THEOREM 5

First we recall some facts about toric geometry, see [3] for details. Let $P \subset \mathbb{R}^d$ be a d -dimensional integral convex polytope. We define a cone

$$\sigma_F = \{v \in \mathbb{R}^d \mid \langle u' - u, v \rangle \geq 0 \ \forall u' \in P, \forall u \in F\}$$

for each face F of P . Then the set

$$\Delta_P = \{\sigma_F \mid F \text{ is a face of } P\}$$

of such cones forms a fan in $\mathbb{R}^d$, which is called the *normal fan* of P . Let $X(\Delta_P)$ be the associated projective toric variety. We denote by $V(\sigma)$ the subvariety of $X(\Delta_P)$ corresponding to $\sigma \in \Delta_P$. Let $\text{Td}_i(X(\Delta_P)) \in A_i(X(\Delta_P))_{\mathbb{Q}}$ be the i -th Todd class in the Chow group of i -cycles with rational coefficients.

Theorem 8. *Let $P \subset \mathbb{R}^d$ be a d -dimensional integral convex polytope and $|(lP) \cap \mathbb{Z}^d| = c_d l^d + c_{d-1} l^{d-1} + \dots + c_0$ be its Ehrhart polynomial. If $\text{Td}_i(X(\Delta_P))$ has an expression of the form $\sum_F r_F [V(\sigma_F)]$ with $r_F \in \mathbb{Q}$, then we have $c_i = \sum_F r_F \text{Vol}(F)$, where $[V(\sigma_F)]$ is the class of $V(\sigma_F)$ in the Chow group and $\text{Vol}(F)$ is the relative volume of F .*

Now we assume that $d = 3$ and P is simple. Then the associated toric variety $X(\Delta_P)$ is $\mathbb{Q}$ -factorial and we know the ring structure of the Chow ring $A^*(X(\Delta_P))_{\mathbb{Q}}$ with rational coefficients. Let $E_1, \dots, E_m$ and $F_1, \dots, F_n$ be the edges and the facets of P , respectively. We have

$$(3.1) \quad \sum_{k=1}^n \langle u, v_k \rangle [V(\sigma_{F_k})] = 0 \quad \forall u \in (\mathbb{Q}^3)^*.$$

If F_{k_1} and F_{k_2} are distinct, then

$$(3.2) \quad [V(\sigma_{F_{k_1}})][V(\sigma_{F_{k_2}})] = \begin{cases} \frac{1}{m(E_j)} [V(\sigma_{E_j})] & (1 \leq \exists j \leq m : F_{k_1} \cap F_{k_2} = E_j), \\ 0 & (F_{k_1} \cap F_{k_2} = \emptyset) \end{cases}$$

in $A^*(X(\Delta_P))_{\mathbb{Q}}$.

Pommersheim gave an expression of $\text{Td}_{d-2}(X(\Delta_P))$ for a d -dimensional simple integral convex polytope $P \subset \mathbb{R}^d$. In the case where $d = 3$, we have the following:

Theorem 9 (Pommersheim [4]). *If $P \subset \mathbb{R}^3$ is a 3-dimensional simple integral convex polytope, then*

$$\text{Td}_1(X(\Delta_P)) = \sum_{j=1}^m \left(s(E_j) + \frac{1}{4} \right) [V(\sigma_{E_j})] + \frac{1}{12} \sum_{k=1}^n [V(\sigma_{F_k})]^2.$$

We use the notation in Definition 3. It suffices to show

$$[V(\sigma_F)]^2 = - \sum_{2 \leq i < j \leq r} a_i \begin{vmatrix} b_{i+1} & \varepsilon_{i+1}^{-1} & 0 & \cdots & 0 \\ \varepsilon_{i+1}^{-1} & b_{i+2} & \varepsilon_{i+2}^{-1} & \ddots & \vdots \\ 0 & \varepsilon_{i+2}^{-1} & \ddots & \ddots & 0 \\ \vdots & \ddots & \ddots & b_{j-2} & \varepsilon_{j-2}^{-1} \\ 0 & \cdots & 0 & \varepsilon_{j-2}^{-1} & b_{j-1} \end{vmatrix} \frac{\varepsilon_i \varepsilon_{i+1} \cdots \varepsilon_{j-1}}{m(P_{j-1} P_j)} [V(\sigma_{P_{j-1} P_j})]$$

for each facet F of P .

We put

$$D(s, t) = \begin{vmatrix} b_s & \varepsilon_s^{-1} & 0 & \cdots & 0 \\ \varepsilon_s^{-1} & b_{s+1} & \varepsilon_{s+1}^{-1} & \ddots & \vdots \\ 0 & \varepsilon_{s+1}^{-1} & \ddots & \ddots & 0 \\ \vdots & \ddots & \ddots & b_{t-1} & \varepsilon_{t-1}^{-1} \\ 0 & \cdots & 0 & \varepsilon_{t-1}^{-1} & b_t \end{vmatrix}$$

for $2 < s \leq t < r$ and $D(s, t) = 1$ for $s > t$. Define $u \in (\mathbb{Q}^3)^*$ by $\langle u, v \rangle = 1, \langle u, v_{k_1} \rangle = 0, \langle u, v_{k_2} \rangle = 0$. By (3.1) and (3.2), we have

$$[V(\sigma_F)]^2 = -[V(\sigma_F)] \sum_{j=1}^r \langle u, v_{k_j} \rangle [V(\sigma_{F_{k_j}})] = - \sum_{j=3}^r \frac{\langle u, v_{k_j} \rangle}{m(P_{j-1} P_j)} [V(\sigma_{P_{j-1} P_j})].$$

Hence it suffices to show

$$(3.3) \quad \langle u, v_{k_j} \rangle = \sum_{i=2}^{j-1} a_i D(i+1, j-1) \varepsilon_i \varepsilon_{i+1} \cdots \varepsilon_{j-1}$$

for any $j = 3, \dots, r$.

First we claim that

$$(3.4) \quad \varepsilon_{j-1}^{-1} v_{k_{j-1}} + \varepsilon_j^{-1} v_{k_{j+1}} = a_j v + b_j v_{k_j}$$

for any $j = 2, \dots, r-1$. By Cramer's rule, we have

$$\begin{aligned} v_{k_{j+1}} &= \frac{\det(v_{k_{j+1}}, v_{k_j}, v_{k_{j-1}})}{\det(v, v_{k_j}, v_{k_{j-1}})} v + \frac{\det(v, v_{k_{j+1}}, v_{k_{j-1}})}{\det(v, v_{k_j}, v_{k_{j-1}})} v_{k_j} + \frac{\det(v, v_{k_j}, v_{k_{j+1}})}{\det(v, v_{k_j}, v_{k_{j-1}})} v_{k_{j-1}} \\ &= \frac{\det(v_{k_{j+1}}, v_{k_j}, v_{k_{j-1}})}{\varepsilon_{j-1}} v + \frac{\det(v, v_{k_{j+1}}, v_{k_{j-1}})}{\varepsilon_{j-1}} v_{k_j} - \frac{\varepsilon_j}{\varepsilon_{j-1}} v_{k_{j-1}}. \end{aligned}$$

So we have

$$(3.5) \quad \begin{aligned} &\varepsilon_{j-1}^{-1} v_{k_{j-1}} + \varepsilon_j^{-1} v_{k_{j+1}} \\ &= \varepsilon_{j-1}^{-1} \varepsilon_j^{-1} \det(v_{k_{j+1}}, v_{k_j}, v_{k_{j-1}}) v + \varepsilon_{j-1}^{-1} \varepsilon_j^{-1} \det(v, v_{k_{j+1}}, v_{k_{j-1}}) v_{k_j}. \end{aligned}$$

Taking the inner product of both sides of (3.5) with $\overrightarrow{P_{j-1}Q_{j-1}}$ gives

$$\varepsilon_j^{-1} \langle \overrightarrow{P_{j-1}Q_{j-1}}, v_{k_{j+1}} \rangle = \varepsilon_{j-1}^{-1} \varepsilon_j^{-1} \det(v_{k_{j+1}}, v_{k_j}, v_{k_{j-1}}) \langle \overrightarrow{P_{j-1}Q_{j-1}}, v \rangle,$$

which means $a_j = \varepsilon_{j-1}^{-1} \varepsilon_j^{-1} \det(v_{k_{j+1}}, v_{k_j}, v_{k_{j-1}})$. Taking the inner product of both sides of (3.5) with $\overrightarrow{P_j P_{j+1}}$ gives

$$\varepsilon_{j-1}^{-1} \langle \overrightarrow{P_j P_{j+1}}, v_{k_{j-1}} \rangle = \varepsilon_{j-1}^{-1} \varepsilon_j^{-1} \det(v, v_{k_{j+1}}, v_{k_{j-1}}) \langle \overrightarrow{P_j P_{j+1}}, v_{k_j} \rangle,$$

which means $b_j = \varepsilon_{j-1}^{-1} \varepsilon_j^{-1} \det(v, v_{k_{j+1}}, v_{k_{j-1}})$. Thus (3.4) follows.

We show (3.3) by induction on j . If $j = 3$, then both sides are $a_2 \varepsilon_2$. If $j = 4$, then both sides are $a_2 b_3 \varepsilon_2 \varepsilon_3 + a_3 \varepsilon_3$. Suppose $4 \leq j \leq r-1$. By (3.4) and the hypothesis of induction, we have

$$\begin{aligned} \langle u, v_{k_{j+1}} \rangle &= \langle u, a_j \varepsilon_j v + b_j \varepsilon_j v_{k_j} - \varepsilon_{j-1}^{-1} \varepsilon_j v_{k_{j-1}} \rangle \\ &= a_j \varepsilon_j + b_j \varepsilon_j \langle u, v_{k_j} \rangle - \varepsilon_{j-1}^{-1} \varepsilon_j \langle u, v_{k_{j-1}} \rangle \\ &= a_j \varepsilon_j + b_j \varepsilon_j \sum_{i=2}^{j-1} a_i D(i+1, j-1) \varepsilon_i \varepsilon_{i+1} \cdots \varepsilon_{j-1} \\ &\quad - \varepsilon_{j-1}^{-1} \varepsilon_j \sum_{i=2}^{j-2} a_i D(i+1, j-2) \varepsilon_i \varepsilon_{i+1} \cdots \varepsilon_{j-2}. \end{aligned}$$

On the other hand,

$$\begin{aligned} &\sum_{i=2}^j a_i D(i+1, j) \varepsilon_i \varepsilon_{i+1} \cdots \varepsilon_j \\ &= a_j \varepsilon_j + a_{j-1} b_j \varepsilon_{j-1} \varepsilon_j + \sum_{i=2}^{j-2} a_i D(i+1, j) \varepsilon_i \varepsilon_{i+1} \cdots \varepsilon_j. \end{aligned}$$

Since

$$\begin{aligned}
& \sum_{i=2}^{j-2} a_i D(i+1, j) \varepsilon_i \varepsilon_{i+1} \cdots \varepsilon_j \\
&= \sum_{i=2}^{j-2} a_i (b_j D(i+1, j-1) - \varepsilon_{j-1}^{-2} D(i+1, j-2)) \varepsilon_i \varepsilon_{i+1} \cdots \varepsilon_j \\
&= b_j \varepsilon_j \sum_{i=2}^{j-2} a_i D(i+1, j-1) \varepsilon_i \varepsilon_{i+1} \cdots \varepsilon_{j-1} \\
&\quad - \varepsilon_{j-1}^{-1} \varepsilon_j \sum_{i=2}^{j-2} a_i D(i+1, j-2) \varepsilon_i \varepsilon_{i+1} \cdots \varepsilon_{j-2},
\end{aligned}$$

we have

$$\begin{aligned}
& \sum_{i=2}^j a_i D(i+1, j) \varepsilon_i \varepsilon_{i+1} \cdots \varepsilon_j \\
&= a_j \varepsilon_j + a_{j-1} b_j \varepsilon_{j-1} \varepsilon_j + b_j \varepsilon_j \sum_{i=2}^{j-2} a_i D(i+1, j-1) \varepsilon_i \varepsilon_{i+1} \cdots \varepsilon_{j-1} \\
&\quad - \varepsilon_{j-1}^{-1} \varepsilon_j \sum_{i=2}^{j-2} a_i D(i+1, j-2) \varepsilon_i \varepsilon_{i+1} \cdots \varepsilon_{j-2} \\
&= a_j \varepsilon_j + b_j \varepsilon_j \sum_{i=2}^{j-1} a_i D(i+1, j-1) \varepsilon_i \varepsilon_{i+1} \cdots \varepsilon_{j-1} \\
&\quad - \varepsilon_{j-1}^{-1} \varepsilon_j \sum_{i=2}^{j-2} a_i D(i+1, j-2) \varepsilon_i \varepsilon_{i+1} \cdots \varepsilon_{j-2} \\
&= \langle u, v_{k_{j+1}} \rangle.
\end{aligned}$$

Thus (3.3) holds for $j+1$. This completes the proof of Theorem 5.

REFERENCES

- [1] M. Beck and S. Robins: Computing the Continuous Discretely, Undergraduate Texts in Mathematics, Springer, 2007.
- [2] E. Ehrhart: Polynômes Arithmétiques et Méthode des Polyèdres en Combinatoire, Birkhäuser, Boston-Basel-Stuttgart, 1977.
- [3] W. Fulton: Introduction to Toric Varieties, Annals of Mathematics Studies **131**, Princeton Univ. Press, Princeton, NJ, 1993.
- [4] J. E. Pommersheim: *Toric varieties, lattice points and Dedekind sums*, Math. Ann. **295** (1993), 1–24.

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA CITY UNIVERSITY,
 3-3-138 SUGIMOTO, SUMIYOSHI-KU, OSAKA 558-8585 JAPAN
E-mail address: d15san0w03@st.osaka-cu.ac.jp